

# Escherichia coli

***Escherichia coli*** (/ˌɛʃəˈrɪkiə ˈkoʊlaɪ/,<sup>[1][2]</sup> also known as ***E. coli*** (/ˌiː ˈkoʊlaɪ/),<sup>[2]</sup> is a Gram-negative, facultative anaerobic, rod-shaped, coliform bacterium of the genus *Escherichia* that is commonly found in the lower intestine of warm-blooded organisms (endotherms).<sup>[3][4]</sup> Most *E. coli* strains are harmless, but some serotypes (EPEC, ETEC etc.) can cause serious food poisoning in their hosts, and are occasionally responsible for food contamination incidents that prompt product recalls.<sup>[5][6]</sup> The harmless strains are part of the normal microbiota of the gut, and can benefit their hosts by producing vitamin K<sub>2</sub>,<sup>[7]</sup> (which helps blood to clot) and preventing colonisation of the intestine with pathogenic bacteria, having a symbiotic relationship.<sup>[8][9]</sup> *E. coli* is expelled into the environment within fecal matter. The bacterium grows massively in fresh fecal matter under aerobic conditions for 3 days, but its numbers decline slowly afterwards.<sup>[10]</sup>

*E. coli* and other facultative anaerobes constitute about 0.1% of gut microbiota,<sup>[11]</sup> and fecal–oral transmission is the major route through which pathogenic strains of the bacterium cause disease. Cells are able to survive outside the body for a limited amount of time, which makes them potential indicator organisms to test environmental samples for fecal contamination.<sup>[12][13]</sup> A growing body of research, though, has examined environmentally persistent *E. coli* which can survive for many days and grow outside a host.<sup>[14]</sup>

The bacterium can be grown and cultured easily and inexpensively in a laboratory setting, and has been intensively investigated for over 60 years. *E. coli* is a chemoheterotroph whose chemically defined medium must include a source of carbon and energy.<sup>[15]</sup> *E. coli* is the most widely studied prokaryotic model organism, and an important species in the fields of biotechnology and microbiology, where it has served as the host organism for the majority of work with recombinant DNA. Under favorable conditions, it takes as little as 20 minutes to reproduce.<sup>[16]</sup>

<i>Escherichia coli</i>
<span></span> <div></div>
Scientific classification <span><span></span></span>
Domain: <span>Bacteria</span>
Phylum: <span>Proteobacteria</span>
Class: <span>Gammaproteobacteria</span>
Order: <span>Enterobacterales</span>
Family: <span>Enterobacteriaceae</span>
Genus: <span><i>Escherichia</i></span>
Species: <span><i><b>E. coli</b></i></span>
Binomial name
<span><i><b>Escherichia coli</b></i></span> <div>(Migula 1895)</div> <div>Castellani and Chalmers 1919</div>
Synonyms
<div> <div><span>■</span> <i>Bacillus coli communis</i></div> <div><span><span>Escherich</span> 1885</span></div> </div>

## Contents

### Biology and biochemistry

- Type and morphology
- Metabolism
- Culture growth
- Cell cycle


Genetic adaptation**Diversity**SerotypesGenome plasticity and evolutionNeotype strainPhylogeny of *E. coli* strains**Genomics****Gene nomenclature****Proteomics**ProteomeInteractome**Normal microbiota**Therapeutic use**Role in disease**Incubation periodTreatmentPrevention**Model organism in life science research**Model organism**History****See also****References****Databases and external links**

## Biology and biochemistry

### Type and morphology

*E. coli* is a Gram-negative, facultative anaerobe (that makes ATP by aerobic respiration if oxygen is present, but is capable of switching to fermentation or anaerobic respiration if oxygen is absent) and nonsporulating bacterium.<sup>[17]</sup> Cells are typically rod-shaped, and are about 2.0  $\mu\text{m}$  long and 0.25–1.0  $\mu\text{m}$  in diameter, with a cell volume of 0.6–0.7  $\mu\text{m}^3$ .<sup>[18][19][20]</sup>

*E. coli* stains Gram-negative because its cell wall is composed of a thin peptidoglycan layer and an outer membrane. During the staining process, *E. coli* picks up the color of the counterstain safranin



Model of successive binary fission in *E. coli*

and stains pink. The outer membrane surrounding the cell wall provides a barrier to certain antibiotics such that *E. coli* is not damaged by penicillin.<sup>[15]</sup>

Strains that possess flagella are motile. The flagella have a peritrichous arrangement.<sup>[21]</sup> It also attaches and effaces to the microvilli of the intestines via an adhesion molecule known as intimin.<sup>[22]</sup>

## Metabolism

*E. coli* can live on a wide variety of substrates and uses mixed acid fermentation in anaerobic conditions, producing lactate, succinate, ethanol, acetate, and carbon dioxide. Since many pathways in mixed-acid fermentation produce hydrogen gas, these pathways require the levels of hydrogen to be low, as is the case when *E. coli* lives together with hydrogen-consuming organisms, such as methanogens or sulphate-reducing bacteria.<sup>[23]</sup>


In addition, *E. coli*'s metabolism can be rewired to solely use CO<sub>2</sub> as the source of carbon for biomass production. In other words, this obligate heterotroph's metabolism can be altered to display autotrophic capabilities by heterologously expressing carbon fixation genes as well as formate dehydrogenase and conducting laboratory evolution experiments. This may be done by using formate to reduce electron carriers and supply the ATP required in anabolic pathways inside of these synthetic autotrophs.<sup>[24]</sup>

## Culture growth

Optimum growth of *E. coli* occurs at 37 °C (98.6 °F), but some laboratory strains can multiply at temperatures up to 49 °C (120 °F).<sup>[25]</sup> *E. coli* grows in a variety of defined laboratory media, such as lysogeny broth, or any medium that contains glucose, ammonium phosphate monobasic, sodium chloride, magnesium sulfate, potassium phosphate dibasic, and water. Growth can be driven by aerobic or anaerobic respiration, using a large variety of redox pairs, including the oxidation of pyruvic acid, formic acid, hydrogen, and amino acids, and the reduction of substrates such as oxygen, nitrate, fumarate, dimethyl sulfoxide, and trimethylamine N-oxide.<sup>[26]</sup> *E. coli* is classified as a facultative anaerobe. It uses oxygen when it is present and available. It can, however, continue to grow in the absence of oxygen using fermentation or anaerobic respiration. The ability to continue growing in the absence of oxygen is an advantage to bacteria because their survival is increased in environments where water predominates.<sup>[15]</sup>

## Cell cycle

The bacterial cell cycle is divided into three stages. The B period occurs between the completion of cell division and the beginning of DNA replication. The C period encompasses the time it takes to replicate the chromosomal DNA. The D period refers to the stage between the conclusion of DNA replication and the end of cell division.<sup>[27]</sup> The doubling rate of *E. coli* is higher when more nutrients are available.



A colony of *E. coli* growing



*E. coli* on sheep blood agar.



*E. coli* growing on basic cultivation media.

However, the length of the C and D periods do not change, even when the doubling time becomes less than the sum of the C and D periods. At the fastest growth rates, replication begins before the previous round of replication has completed, resulting in multiple replication forks along the DNA and overlapping cell cycles.<sup>[28]</sup>

The number of replication forks in fast growing *E. coli* typically follows  $2n$  ( $n = 1, 2$  or  $3$ ). This only happens if replication is initiated simultaneously from all origins of replications, and is referred to as synchronous replication. However, not all cells in a culture replicate synchronously. In this case cells do not have multiples of two replication forks. Replication initiation is then referred to being asynchronous.<sup>[29]</sup> However, asynchrony can be caused by mutations to for instance DnaA<sup>[29]</sup> or DnaA initiator-associating protein DiaA.<sup>[30]</sup>

## Genetic adaptation

*E. coli* and related bacteria possess the ability to transfer DNA via bacterial conjugation or transduction, which allows genetic material to spread horizontally through an existing population. The process of transduction, which uses the bacterial virus called a bacteriophage,<sup>[31]</sup> is where the spread of the gene encoding for the Shiga toxin from the Shigella bacteria to *E. coli* helped produce *E. coli* O157:H7, the Shiga toxin-producing strain of *E. coli*.

## Diversity

*E. coli* encompasses an enormous population of bacteria that exhibit a very high degree of both genetic and phenotypic diversity. Genome sequencing of many isolates of *E. coli* and related bacteria shows that a taxonomic reclassification would be desirable. However, this has not been done, largely due to its medical importance,<sup>[32]</sup> and *E. coli* remains one of the most diverse bacterial species: only 20% of the genes in a typical *E. coli* genome is shared among all strains.<sup>[33]</sup>

In fact, from the more constructive point of view, the members of genus *Shigella* (*S. dysenteriae*, *S. flexneri*, *S. boydii*, and *S. sonnei*) should be classified as *E. coli* strains, a phenomenon termed taxa in disguise.<sup>[34]</sup> Similarly, other strains of *E. coli* (e.g. the K-12 strain commonly used in recombinant DNA work) are sufficiently different that they would merit reclassification.



Scanning electron micrograph of an *E. coli* colony.

A strain is a subgroup within the species that has unique characteristics that distinguish it from other strains. These differences are often detectable only at the molecular level; however, they may result in changes to the physiology or lifecycle of the bacterium. For example, a strain may gain pathogenic capacity, the ability to use a unique carbon source, the ability to take upon a particular ecological niche, or the ability to resist antimicrobial agents. Different strains of *E. coli* are often host-specific, making it possible to determine the source of fecal contamination in environmental samples.<sup>[12][13]</sup> For example, knowing which *E. coli* strains are present in a water sample allows researchers to make assumptions about whether the contamination originated from a human, another mammal, or a bird.

## Serotypes



A common subdivision system of *E. coli*, but not based on evolutionary relatedness, is by serotype, which is based on major surface antigens (O antigen: part of lipopolysaccharide layer; H: flagellin; K antigen: capsule), e.g. O157:H7).<sup>[35]</sup> It is, however, common to cite only the serogroup, i.e. the O-antigen. At present, about 190 serogroups are known.<sup>[36]</sup> The common laboratory strain has a mutation that prevents the formation of an O-antigen and is thus not typeable.

## Genome plasticity and evolution

Like all lifeforms, new strains of *E. coli* evolve through the natural biological processes of mutation, gene duplication, and horizontal gene transfer; in particular, 18% of the genome of the laboratory strain MG1655 was horizontally acquired since the divergence from *Salmonella*.<sup>[37]</sup> *E. coli* K-12 and *E. coli* B strains are the most frequently used varieties for laboratory purposes. Some strains develop traits that can be harmful to a host animal. These virulent strains typically cause a bout of diarrhea that is often self-limiting in healthy adults but is frequently lethal to children in the developing world.<sup>[38]</sup> More virulent strains, such as O157:H7, cause serious illness or death in the elderly, the very young, or the immunocompromised.<sup>[38][39]</sup>

The genera *Escherichia* and *Salmonella* diverged around 102 million years ago (credibility interval: 57–176 mya), which coincides with the divergence of their hosts: the former being found in mammals and the latter in birds and reptiles.<sup>[40]</sup> This was followed by a split of an *Escherichia* ancestor into five species (*E. albertii*, *E. coli*, *E. fergusonii*, *E. hermannii*, and *E. vulneris*). The last *E. coli* ancestor split between 20 and 30 million years ago.<sup>[41]</sup>

The long-term evolution experiments using *E. coli*, begun by Richard Lenski in 1988, have allowed direct observation of genome evolution over more than 65,000 generations in the laboratory.<sup>[42]</sup> For instance, *E. coli* typically do not have the ability to grow aerobically with citrate as a carbon source, which is used as a diagnostic criterion with which to differentiate *E. coli* from other, closely, related bacteria such as *Salmonella*. In this experiment, one population of *E. coli* unexpectedly evolved the ability to aerobically metabolize citrate, a major evolutionary shift with some hallmarks of microbial speciation.

In the microbial world, a relationship of predation can be established similar to that observed in the animal world. Considered, it has been seen that *E. coli* is the prey of multiple generalist predators, such as *Myxococcus xanthus*. In this predator-prey relationship, a parallel evolution of both species is observed through genomic and phenotypic modifications, in the case of *E. coli* the modifications are modified in two aspects involved in their virulence such as mucoid production (excessive production of exoplasmic acid alginate ) and the suppression of the OmpT gene, producing in future generations a better adaptation of one of the species that is counteracted by the evolution of the other, following a co-evolutionary model demonstrated by the Red Queen hypothesis.<sup>[43]</sup>

## Neotype strain

*E. coli* is the type species of the genus (*Escherichia*) and in turn *Escherichia* is the type genus of the family Enterobacteriaceae, where the family name does not stem from the genus *Enterobacter* + "i" (sic.) + "aceae", but from "enterobacterium" + "aceae" (enterobacterium being not a genus, but an alternative trivial name to enteric bacterium).<sup>[44][45][46]</sup>


The original strain described by Escherich is believed to be lost, consequently a new type strain (neotype) was chosen as a representative: the neotype strain is U5/41<sup>T</sup>,<sup>[47]</sup> also known under the deposit names DSM 30083,<sup>[48]</sup> ATCC 11775,<sup>[49]</sup> and NCTC 9001,<sup>[50]</sup> which is pathogenic to chickens and has an


O1:K1:H7 serotype.<sup>[51]</sup> However, in most studies, either O157:H7, K-12 MG1655, or K-12 W3110 were used as a representative *E. coli*. The genome of the type strain has only lately been sequenced.<sup>[47]</sup>


## Phylogeny of *E. coli* strains

Many strains belonging to this species have been isolated and characterised. In addition to serotype (*vide supra*), they can be classified according to their phylogeny, i.e. the inferred evolutionary history, as shown below where the species is divided into six groups.<sup>[52][53]</sup> Particularly the use of whole genome sequences yields highly supported phylogenies. Based on such data, five subspecies of *E. coli* were distinguished.<sup>[47]</sup>

The link between phylogenetic distance ("relatedness") and pathology is small,<sup>[47]</sup> e.g. the O157:H7 serotype strains, which form a clade ("an exclusive group")—group E below—are all enterohaemorrhagic strains (EHEC), but not all EHEC strains are closely related. In fact, four different species of *Shigella* are nested among *E. coli* strains (*vide supra*), while *E. albertii* and *E. fergusonii* are outside this group. Indeed, all *Shigella* species were placed within a single subspecies of *E. coli* in a phylogenomic study that included the type strain,<sup>[47]</sup> and for this reason an according reclassification is difficult. All commonly used research strains of *E. coli* belong to group A and are derived mainly from Clifton's K-12 strain ( $\lambda^+$  F<sup>+</sup>; O16) and to a lesser degree from d'Herelle's *Bacillus coli* strain (B strain)(O7).









*E. coli* BL21-DE3 (O7, expression molecular biology strain with T7 polymerase for pET system)

## Genomics

The first complete DNA sequence of an *E. coli* genome (laboratory strain K-12 derivative MG1655) was published in 1997. It is a circular DNA molecule 4.6 million base pairs in length, containing 4288 annotated protein-coding genes (organized into 2584 operons), seven ribosomal RNA (rRNA) operons, and 86 transfer RNA (tRNA) genes. Despite having been the subject of intensive genetic analysis for about 40 years, many of these genes were previously unknown. The coding density was found to be very high, with a mean distance between genes of only 118 base pairs. The genome was observed to contain a significant number of transposable genetic elements, repeat elements, cryptic prophages, and bacteriophage remnants.<sup>[54]</sup>

More than three hundred complete genomic sequences of *Escherichia* and *Shigella* species are known. The genome sequence of the type strain of *E. coli* was added to this collection before 2014.<sup>[47]</sup> Comparison of these sequences shows a remarkable amount of diversity; only about 20% of each genome represents sequences present in every one of the isolates, while around 80% of each genome can vary among isolates.<sup>[33]</sup> Each individual genome contains between 4,000 and 5,500 genes, but the total number of different genes among all of the sequenced *E. coli* strains (the pangenome) exceeds 16,000. This very large variety of component genes has been interpreted to mean that two-thirds of the *E. coli* pangenome originated in other species and arrived through the process of horizontal gene transfer.<sup>[55]</sup>



An image of *E. coli* using early electron microscopy.

## Gene nomenclature

Genes in *E. coli* are usually named by 4-letter acronyms that derive from their function (when known) and italicized. For instance, *recA* is named after its role in homologous recombination plus the letter A. Functionally related genes are named *recB*, *recC*, *recD* etc. The proteins are named by uppercase acronyms, e.g. RecA, RecB, etc. When the genome of *E. coli* was sequenced, all genes were numbered (more or less) in their order on the genome and abbreviated by b numbers, such as b2819 (= *recD*). The "b" names were created after Fred Blattner, who led the genome sequence effort.<sup>[54]</sup> Another numbering system was introduced with the sequence of another *E. coli* strain, W3110, which was sequenced in Japan and hence uses numbers starting by JW... (Japanese W3110), e.g. JW2787 (= *recD*).<sup>[56]</sup> Hence, *recD* = b2819 = JW2787. Note, however, that most databases have their own numbering system, e.g. the EcoGene database<sup>[57]</sup> uses EG10826 for *recD*. Finally, ECK numbers are specifically used for alleles in the MG1655 strain of *E. coli* K-12.<sup>[57]</sup> Complete lists of genes and their synonyms can be obtained from databases such as EcoGene or Uniprot.

## Proteomics

## Proteome

Several studies have investigated the proteome of *E. coli*. By 2006, 1,627 (38%) of the 4,237 open reading frames (ORFs) had been identified experimentally.<sup>[58]</sup> The 4,639,221–base pair sequence of *Escherichia coli* K-12 is presented. Of 4288 protein-coding genes annotated, 38 percent have no attributed function. Comparison with five other sequenced microbes reveals ubiquitous as well as narrowly distributed gene families; many families of similar genes within *E. coli* are also evident. The largest family of paralogous proteins contains 80 ABC transporters. The genome as a whole is strikingly organized with respect to the local direction of replication; guanines, oligonucleotides possibly related to replication and recombination, and most genes are so oriented. The genome also contains insertion sequence (IS) elements, phage remnants, and many other patches of unusual composition indicating genome plasticity through horizontal transfer.<sup>[54]</sup>

## Interactome

The interactome of *E. coli* has been studied by affinity purification and mass spectrometry (AP/MS) and by analyzing the binary interactions among its proteins.

**Protein complexes.** A 2006 study purified 4,339 proteins from cultures of strain K-12 and found interacting partners for 2,667 proteins, many of which had unknown functions at the time.<sup>[59]</sup> A 2009 study found 5,993 interactions between proteins of the same *E. coli* strain, though these data showed little overlap with those of the 2006 publication.<sup>[60]</sup>

**Binary interactions.** Rajagopala *et al.* (2014) have carried out systematic yeast two-hybrid screens with most *E. coli* proteins, and found a total of 2,234 protein-protein interactions.<sup>[61]</sup> This study also integrated genetic interactions and protein structures and mapped 458 interactions within 227 protein complexes.

## Normal microbiota

---

*E. coli* belongs to a group of bacteria informally known as coliforms that are found in the gastrointestinal tract of warm-blooded animals.<sup>[44]</sup> *E. coli* normally colonizes an infant's gastrointestinal tract within 40 hours of birth, arriving with food or water or from the individuals handling the child. In the bowel, *E. coli* adheres to the mucus of the large intestine. It is the primary facultative anaerobe of the human gastrointestinal tract.<sup>[62]</sup> (Facultative anaerobes are organisms that can grow in either the presence or absence of oxygen.) As long as these bacteria do not acquire genetic elements encoding for virulence factors, they remain benign commensals.<sup>[63]</sup>

## Therapeutic use

Nonpathogenic *E. coli* strain Nissle 1917, (Mutaflor) and *E. coli* O83:K24:H31 (Colinfant)<sup>[64][65]</sup> are used as probiotic agents in medicine, mainly for the treatment of various gastrointestinal diseases,<sup>[66]</sup> including inflammatory bowel disease.<sup>[67]</sup>

## Role in disease

---

Most *E. coli* strains do not cause disease, naturally living in the gut,<sup>[68]</sup> but virulent strains can cause gastroenteritis, urinary tract infections, neonatal meningitis, hemorrhagic colitis, and Crohn's disease. Common signs and symptoms include severe abdominal cramps, diarrhea, hemorrhagic colitis, vomiting, and sometimes fever. In rarer cases, virulent strains are also responsible for bowel necrosis (tissue death) and perforation without progressing to hemolytic-uremic syndrome, peritonitis, mastitis, sepsis, and Gram-negative pneumonia. Very young children are more susceptible to develop severe illness, such as hemolytic uremic syndrome; however, healthy individuals of all ages are at risk to the severe consequences that may arise as a result of being infected with *E. coli*.<sup>[62][69][70][71]</sup>

Some strains of *E. coli*, for example O157:H7, can produce Shiga toxin (classified as a bioterrorism agent). The Shiga toxin causes inflammatory responses in target cells of the gut, leaving behind lesions which result in the bloody diarrhea that is a symptom of a Shiga toxin-producing *E. coli* (STEC) infection. This toxin further causes premature destruction of the red blood cells, which then clog the body's filtering system, the kidneys, in some rare cases (usually in children and the elderly) causing hemolytic-uremic syndrome (HUS), which may lead to kidney failure and even death. Signs of hemolytic uremic syndrome include decreased frequency of urination, lethargy, and paleness of cheeks and inside the lower eyelids. In 25% of HUS patients, complications of nervous system occur, which in turn causes strokes. In addition, this strain causes the buildup of fluid (since the kidneys do not work), leading to edema around the lungs and legs and arms. This increase in fluid buildup especially around the lungs impedes the functioning of the heart, causing an increase in blood pressure.<sup>[72][22][73][74][75][70][71]</sup>

Uropathogenic *E. coli* (UPEC) is one of the main causes of urinary tract infections.<sup>[76]</sup> It is part of the normal microbiota in the gut and can be introduced in many ways. In particular for females, the direction of wiping after defecation (wiping back to front) can lead to fecal contamination of the urogenital orifices. Anal intercourse can also introduce this bacterium into the male urethra, and in switching from anal to vaginal intercourse, the male can also introduce UPEC to the female urogenital system.

Enterotoxigenic *E. coli* (ETEC) is the most common cause of traveler's diarrhea, with as many as 840 million cases worldwide in developing countries each year. The bacteria, typically transmitted through contaminated food or drinking water, adheres to the intestinal lining, where it secretes either of two types of enterotoxins, leading to watery diarrhea. The rate and severity of infections are higher among children under the age of five, including as many as 380,000 deaths annually.<sup>[77]</sup>

In May 2011, one *E. coli* strain, O104:H4, was the subject of a bacterial outbreak that began in Germany. Certain strains of *E. coli* are a major cause of foodborne illness. The outbreak started when several people in Germany were infected with enterohemorrhagic *E. coli* (EHEC) bacteria, leading to hemolytic-uremic syndrome (HUS), a medical emergency that requires urgent treatment. The outbreak did not only concern Germany, but also 15 other countries, including regions in North America.<sup>[78]</sup> On 30 June 2011, the German *Bundesinstitut für Risikobewertung* (BfR) (Federal Institute for Risk Assessment, a federal institute within the German Federal Ministry of Food, Agriculture and Consumer Protection) announced that seeds of fenugreek from Egypt were likely the cause of the EHEC outbreak.<sup>[79]</sup>

## Incubation period

The time between ingesting the STEC bacteria and feeling sick is called the "incubation period". The incubation period is usually 3–4 days after the exposure, but may be as short as 1 day or as long as 10 days. The symptoms often begin slowly with mild belly pain or non-bloody diarrhea that worsens over several days. HUS, if it occurs, develops an average 7 days after the first symptoms, when the diarrhea is improving.<sup>[80]</sup>

## Treatment

The mainstay of treatment is the assessment of dehydration and replacement of fluid and electrolytes. Administration of antibiotics has been shown to shorten the course of illness and duration of excretion of enterotoxigenic *E. coli* (ETEC) in adults in endemic areas and in traveller's diarrhea, though the rate of resistance to commonly used antibiotics is increasing and they are generally not recommended.<sup>[81]</sup> The antibiotic used depends upon susceptibility patterns in the particular geographical region. Currently, the antibiotics of choice are fluoroquinolones or azithromycin, with an emerging role for rifaximin. Oral rifaximin, a semisynthetic rifamycin derivative, is an effective and well-tolerated antibacterial for the management of adults with non-invasive traveller's diarrhea. Rifaximin was significantly more effective than placebo and no less effective than ciprofloxacin in reducing the duration of diarrhea. While rifaximin is effective in patients with *E. coli*-predominant traveller's diarrhea, it appears ineffective in patients infected with inflammatory or invasive enteropathogens.<sup>[82]</sup>

## Prevention

ETEC is the type of *E. coli* that most vaccine development efforts are focused on. Antibodies against the LT and major CFs of ETEC provide protection against LT-producing, ETEC-expressing homologous CFs. Oral inactivated vaccines consisting of toxin antigen and whole cells, i.e. the licensed recombinant cholera B subunit (rCTB)-WC cholera vaccine Dukoral, have been developed. There are currently no licensed vaccines for ETEC, though several are in various stages of development.<sup>[83]</sup> In different trials, the rCTB-WC cholera vaccine provided high (85–100%) short-term protection. An oral ETEC vaccine candidate consisting of rCTB and formalin inactivated *E. coli* bacteria expressing major CFs has been shown in clinical trials to be safe, immunogenic, and effective against severe diarrhoea in American travelers but not against ETEC diarrhoea in young children in Egypt. A modified ETEC vaccine consisting of recombinant *E. coli* strains over-expressing the major CFs and a more LT-like hybrid toxoid called LCTBA, are undergoing clinical testing.<sup>[84]</sup> <sup>[85]</sup>

Other proven prevention methods for *E. coli* transmission include handwashing and improved sanitation and drinking water, as transmission occurs through fecal contamination of food and water supplies. Additionally, thoroughly cooking meat and avoiding consumption of raw, unpasteurized beverages, such as juices and milk are other proven methods for preventing *E.coli*. Lastly, avoid cross-contamination of utensils and work spaces when preparing food.<sup>[86]</sup>

## Model organism in life science research

---

Because of its long history of laboratory culture and ease of manipulation, *E. coli* plays an important role in modern biological engineering and industrial microbiology.<sup>[87]</sup> The work of Stanley Norman Cohen and Herbert Boyer in *E. coli*, using plasmids and restriction enzymes to create recombinant DNA, became a foundation of biotechnology.<sup>[88]</sup>

*E. coli* is a very versatile host for the production of heterologous proteins,<sup>[89]</sup> and various protein expression systems have been developed which allow the production of recombinant proteins in *E. coli*. Researchers can introduce genes into the microbes using plasmids which permit high level expression of protein, and such protein may be mass-produced in industrial fermentation processes. One of the first useful applications of recombinant DNA technology was the manipulation of *E. coli* to produce human insulin.<sup>[90]</sup>



Many proteins previously thought difficult or impossible to be expressed in *E. coli* in folded form have been successfully expressed in *E. coli*. For example, proteins with multiple disulphide bonds may be produced in the periplasmic space or in the cytoplasm of mutants rendered sufficiently oxidizing to allow disulphide-bonds to form,<sup>[91]</sup> while proteins requiring post-translational modification such as glycosylation for stability or function have been expressed using the N-linked glycosylation system of *Campylobacter jejuni* engineered into *E. coli*.<sup>[92][93][94]</sup>

Modified *E. coli* cells have been used in vaccine development, bioremediation, production of biofuels,<sup>[95]</sup> lighting, and production of immobilised enzymes.<sup>[89][96]</sup>

Strain K-12 is a mutant form of *E. coli* that over-expresses the enzyme Alkaline Phosphatase (ALP).<sup>[97]</sup> The mutation arises due to a defect in the gene that constantly codes for the enzyme. A gene that is producing a product without any inhibition is said to have constitutive activity. This particular mutant form is used to isolate and purify the aforementioned enzyme.<sup>[97]</sup>

Strain OP50 of *Escherichia coli* is used for maintenance of *Caenorhabditis elegans* cultures.

Strain JM109 is a mutant form of *E. coli* that is *recA* and *endA* deficient. The strain can be utilized for blue/white screening when the cells carry the fertility factor episome<sup>[98]</sup> Lack of *recA* decreases the possibility of unwanted restriction of the DNA of interest and lack of *endA* inhibit plasmid DNA decomposition. Thus, JM109 is useful for cloning and expression systems.

## Model organism

*E. coli* is frequently used as a model organism in microbiology studies. Cultivated strains (e.g. *E. coli* K12) are well-adapted to the laboratory environment, and, unlike wild-type strains, have lost their ability to thrive in the intestine. Many laboratory strains lose their ability to form biofilms.<sup>[99][100]</sup> These features protect wild-type strains from antibodies and other chemical attacks, but require a large expenditure of energy and material resources. *E. coli* is often used as a representative microorganism in the research of novel water treatment and sterilisation methods, including photocatalysis. By standard plate count methods, following sequential dilutions, and growth on agar gel plates, the concentration of viable organisms or CFUs (Colony Forming Units), in a known volume of treated water can be evaluated, allowing the comparative assessment of materials performance.<sup>[101]</sup>

In 1946, Joshua Lederberg and Edward Tatum first described the phenomenon known as bacterial conjugation using *E. coli* as a model bacterium,<sup>[102]</sup> and it remains the primary model to study conjugation.<sup>[103]</sup> *E. coli* was an integral part of the first experiments to understand phage genetics,<sup>[104]</sup> and early researchers, such as Seymour Benzer, used *E. coli* and phage T4 to understand the topography of gene structure.<sup>[105]</sup> Prior to Benzer's research, it was not known whether the gene was a linear structure, or if it had a branching pattern.<sup>[106]</sup>

*E. coli* was one of the first organisms to have its genome sequenced; the complete genome of *E. coli* K12 was published by *Science* in 1997<sup>[54]</sup>

From 2002 to 2010, a team at the Hungarian Academy of Science created a strain of *Escherichia coli* called MDS42, which is now sold by Scarab Genomics of Madison, WI under the name of "Clean Genome. E.coli",<sup>[107]</sup> where 15% of the genome of the parental strain (*E. coli* K-12 MG1655) were removed to aid in molecular biology efficiency, removing IS elements, pseudogenes and phages, resulting in better maintenance of plasmid-encoded toxic genes, which are often inactivated by transposons.<sup>[108][109][110]</sup> Biochemistry and replication machinery were not altered.

By evaluating the possible combination of nanotechnologies with landscape ecology, complex habitat landscapes can be generated with details at the nanoscale.<sup>[111]</sup> On such synthetic ecosystems, evolutionary experiments with *E. coli* have been performed to study the spatial biophysics of adaptation in an island biogeography on-chip.

Studies are also being performed attempting to program *E. coli* to solve complicated mathematics problems, such as the Hamiltonian path problem.<sup>[112]</sup>

In other studies, non-pathogenic *E. coli* has been used as a model microorganism towards understanding the effects of simulated microgravity (on Earth) on the same.<sup>[113][114]</sup>

## History

In 1885, the German-Austrian pediatrician Theodor Escherich discovered this organism in the feces of healthy individuals. He called it *Bacterium coli commune* because it is found in the colon. Early classifications of prokaryotes placed these in a handful of genera based on their shape and motility (at that time Ernst Haeckel's classification of bacteria in the kingdom Monera was in place).<sup>[85][115][116]</sup>

*Bacterium coli* was the type species of the now invalid genus *Bacterium* when it was revealed that the former type species ("*Bacterium trilolare*") was missing.<sup>[117]</sup> Following a revision of *Bacterium*, it was reclassified as *Bacillus coli* by Migula in 1895<sup>[118]</sup> and later reclassified in the newly created genus *Escherichia*, named after its original discoverer.<sup>[119]</sup>

In 1996 the world's worst outbreak of *E. coli* food poisoning occurred in Wishaw, Scotland, killing 21 people.<sup>[120]</sup>

## See also

- Bacteriological water analysis
- Carbon monoxide-releasing molecules
- Contamination control
- Dam dcm strain
- Eijkman test
- Fecal coliform
- International Code of Nomenclature of Bacteria
- List of strains of *Escherichia coli*
- Mannan oligosaccharide-based nutritional supplements
- Overflow metabolism
- T4 rII system

## References

- "coli" (<http://oed.com/search?searchType=dictionary&q=coli>). *Oxford English Dictionary* (3rd ed.). Oxford University Press. September 2005. (Subscription or UK public library membership (<https://www.oed.com/public/login/loggingin#withyourlibrary>) required.)
- Wells, J. C. (2000) Longman Pronunciation Dictionary. Harlow [England], Pearson Education Ltd.
- Tenaillon O, Skurnik D, Picard B, Denamur E (March 2010). "The population genetics of commensal *Escherichia coli*". *Nature Reviews. Microbiology*. **8** (3): 207–17. doi:10.1038/nrmicro2298 (<https://doi.org/10.1038/nrmicro2298>). PMID 20157339 (<https://pubmed.ncbi.nlm.nih.gov/20157339>). S2CID 5490303 (<https://api.semanticscholar.org/CorpusID:5490303>).
- Singleton P (1999). *Bacteria in Biology, Biotechnology and Medicine* (5th ed.). Wiley. pp. 444–54. ISBN 978-0-471-98880-9.

5. "Escherichia coli" (<https://www.cdc.gov/ecoli/index.html/>). *CDC National Center for Emerging and Zoonotic Infectious Diseases*. Retrieved 2 October 2012.
6. Vogt RL, Dippold L (2005). "Escherichia coli O157:H7 outbreak associated with consumption of ground beef, June–July 2002" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1497708>). *Public Health Reports*. **120** (2): 174–78. doi:10.1177/003335490512000211 (<https://doi.org/10.1177%2F003335490512000211>). PMC 1497708 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1497708>). PMID 15842119 (<https://pubmed.ncbi.nlm.nih.gov/15842119>).
7. Bentley R, Meganathan R (September 1982). "Biosynthesis of vitamin K (menaquinone) in bacteria" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC281544>). *Microbiological Reviews*. **46** (3): 241–80. doi:10.1128/MMBR.46.3.241-280.1982 (<https://doi.org/10.1128%2FMMBR.46.3.241-280.1982>). PMC 281544 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC281544>). PMID 6127606 (<https://pubmed.ncbi.nlm.nih.gov/6127606>).
8. Hudault S, Guignot J, Servin AL (July 2001). "Escherichia coli strains colonising the gastrointestinal tract protect germfree mice against Salmonella typhimurium infection" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1728375>). *Gut*. **49** (1): 47–55. doi:10.1136/gut.49.1.47 (<https://doi.org/10.1136%2Fgut.49.1.47>). PMC 1728375 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1728375>). PMID 11413110 (<https://pubmed.ncbi.nlm.nih.gov/11413110>).
9. Reid G, Howard J, Gan BS (September 2001). "Can bacterial interference prevent infection?". *Trends in Microbiology*. **9** (9): 424–28. doi:10.1016/S0966-842X(01)02132-1 (<https://doi.org/10.1016%2FS0966-842X%2801%2902132-1>). PMID 11553454 (<https://pubmed.ncbi.nlm.nih.gov/11553454>).
10. Russell JB, Jarvis GN (April 2001). "Practical mechanisms for interrupting the oral-fecal lifecycle of Escherichia coli". *Journal of Molecular Microbiology and Biotechnology*. **3** (2): 265–72. PMID 11321582 (<https://pubmed.ncbi.nlm.nih.gov/11321582>).
11. Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, et al. (June 2005). "Diversity of the human intestinal microbial flora" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1395357>). *Science*. **308** (5728): 1635–38. Bibcode:2005Sci...308.1635E (<https://ui.adsabs.harvard.edu/abs/2005Sci...308.1635E>). doi:10.1126/science.1110591 (<https://doi.org/10.1126%2Fscience.1110591>). PMC 1395357 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1395357>). PMID 15831718 (<https://pubmed.ncbi.nlm.nih.gov/15831718>).
12. Feng P; Weagant S; Grant, M (1 September 2002). "Enumeration of *Escherichia coli* and the Coliform Bacteria" (<https://web.archive.org/web/20090519200935/http://www.cfsan.fda.gov/~ebam/bam-4.html>). *Bacteriological Analytical Manual* (8th ed.). FDA/Center for Food Safety & Applied Nutrition. Archived from the original (<http://www.cfsan.fda.gov/~ebam/bam-4.html>) on 19 May 2009. Retrieved 25 January 2007.
13. Thompson A (4 June 2007). "E. coli Thrives in Beach Sands" ([http://www.livescience.com/health/070604\\_beach\\_ecoli.html](http://www.livescience.com/health/070604_beach_ecoli.html)). Live Science. Retrieved 3 December 2007.
14. Montealegre MC, Roy S, Böni F, Hossain MI, Navab-Daneshmand T, Caduff L, et al. (December 2018). "Risk Factors for Detection, Survival, and Growth of Antibiotic-Resistant and Pathogenic Escherichia coli in Household Soils in Rural Bangladesh" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6275341>). *Applied and Environmental Microbiology*. **84** (24): e01978–18. doi:10.1128/AEM.01978-18 (<https://doi.org/10.1128%2FAEM.01978-18>). PMC 6275341 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6275341>). PMID 30315075 (<https://pubmed.ncbi.nlm.nih.gov/30315075>).
15. Tortora, Gerard (2010). *Microbiology: An Introduction*. San Francisco, CA: Benjamin Cummings. pp. 85–87, 161, 165. ISBN 978-0-321-55007-1.
16. "Bacteria" (<http://www.microbiologyonline.org.uk/about-microbiology/introducing-microbes/bacteria>). Microbiologyonline. Archived (<https://web.archive.org/web/20140227212658/http://www.microbiologyonline.org.uk/about-microbiology/introducing-microbes/bacteria>) from the original on 27 February 2014. Retrieved 27 February 2014.

17. "E.Coli" ([http://www.redorbit.com/education/reference\\_library/health\\_1/bacteria/2584144/escherichia\\_coli/](http://www.redorbit.com/education/reference_library/health_1/bacteria/2584144/escherichia_coli/)). Redorbit. Retrieved 27 November 2013.
18. "Facts about E. coli: dimensions, as discussed in bacteria: Diversity of structure of bacteria" (<http://www.britannica.com/science/bacteria/Diversity-of-structure-of-bacteria>). *Britannica.com – Britannica Online Encyclopedia*. Retrieved 25 June 2015.
19. Yu AC, Loo JF, Yu S, Kong SK, Chan TF (January 2014). "Monitoring bacterial growth using tunable resistive pulse sensing with a pore-based technique". *Applied Microbiology and Biotechnology*. **98** (2): 855–62. doi:10.1007/s00253-013-5377-9 (<https://doi.org/10.1007/s00253-013-5377-9>). PMID 24287933 (<https://pubmed.ncbi.nlm.nih.gov/24287933>). S2CID 2956197 (<https://api.semanticscholar.org/CorpusID:2956197>).
20. Kubitschek HE (January 1990). "Cell volume increase in Escherichia coli after shifts to richer media" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC208405>). *Journal of Bacteriology*. **172** (1): 94–101. doi:10.1128/jb.172.1.94-101.1990 (<https://doi.org/10.1128/jb.172.1.94-101.1990>). PMC 208405 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC208405>). PMID 2403552 (<https://pubmed.ncbi.nlm.nih.gov/2403552>).
21. Darnton NC, Turner L, Rojevsky S, Berg HC (March 2007). "On torque and tumbling in swimming Escherichia coli" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1855780>). *Journal of Bacteriology*. **189** (5): 1756–64. doi:10.1128/JB.01501-06 (<https://doi.org/10.1128/JB.01501-06>). PMC 1855780 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1855780>). PMID 17189361 (<https://pubmed.ncbi.nlm.nih.gov/17189361>).
22. "E. Coli O157 in North America – microbewiki" ([https://microbewiki.kenyon.edu/index.php/E.\\_coli\\_O157\\_in\\_North\\_America](https://microbewiki.kenyon.edu/index.php/E._coli_O157_in_North_America)).
23. Madigan MT, Martinko JM (2006). *Brock Biology of microorganisms* (11th ed.). Pearson. ISBN 978-0-13-196893-6.
24. Gleizer S, Ben-Nissan R, Bar-On YM, Antonovsky N, Noor E, Zohar Y, et al. (November 2019). "2" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6904909>). *Cell*. **179** (6): 1255–1263.e12. doi:10.1016/j.cell.2019.11.009 (<https://doi.org/10.1016/j.cell.2019.11.009>). PMC 6904909 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6904909>). PMID 31778652 (<https://pubmed.ncbi.nlm.nih.gov/31778652>).
25. Fotadar U, Zaveloff P, Terracio L (2005). "Growth of Escherichia coli at elevated temperatures". *Journal of Basic Microbiology*. **45** (5): 403–04. doi:10.1002/jobm.200410542 (<https://doi.org/10.1002/jobm.200410542>). PMID 16187264 (<https://pubmed.ncbi.nlm.nih.gov/16187264>).
26. Ingledew WJ, Poole RK (September 1984). "The respiratory chains of Escherichia coli" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC373010>). *Microbiological Reviews*. **48** (3): 222–71. doi:10.1128/MMBR.48.3.222-271.1984 (<https://doi.org/10.1128/MMBR.48.3.222-271.1984>). PMC 373010 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC373010>). PMID 6387427 (<https://pubmed.ncbi.nlm.nih.gov/6387427>).
27. Wang JD, Levin PA (November 2009). "Metabolism, cell growth and the bacterial cell cycle" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2887316>). *Nature Reviews. Microbiology*. **7** (11): 822–27. doi:10.1038/nrmicro2202 (<https://doi.org/10.1038/nrmicro2202>). PMC 2887316 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2887316>). PMID 19806155 (<https://pubmed.ncbi.nlm.nih.gov/19806155>).
28. Cooper S, Helmstetter CE (February 1968). "Chromosome replication and the division cycle of Escherichia coli B/r". *Journal of Molecular Biology*. **31** (3): 519–40. doi:10.1016/0022-2836(68)90425-7 ([https://doi.org/10.1016/0022-2836\(68\)90425-7](https://doi.org/10.1016/0022-2836(68)90425-7)). PMID 4866337 (<https://pubmed.ncbi.nlm.nih.gov/4866337>).



29. Skarstad, K.; Boye, E.; Steen, H.b. (1 July 1986). "Timing of initiation of chromosome replication in individual *Escherichia coli* cells" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1166998>). *The EMBO Journal*. **5** (7): 1711–17. doi:10.1002/j.1460-2075.1986.tb04415.x (<https://doi.org/10.1002%2Fj.1460-2075.1986.tb04415.x>). ISSN 0261-4189 (<https://www.worldcat.org/issn/0261-4189>). PMC 1166998 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1166998>). PMID 3527695 (<https://pubmed.ncbi.nlm.nih.gov/3527695>).
30. Ishida, Takuma; Akimitsu, Nobuyoshi; Kashioka, Tamami; Hatano, Masakazu; Kubota, Toshio; Ogata, Yasuyuki; Sekimizu, Kazuhisa; Katayama, Tsutomu (29 October 2004). "DiaA, a Novel DnaA-binding Protein, Ensures the Timely Initiation of *Escherichia coli* Chromosome Replication" (<https://doi.org/10.1074/jbc.M402762200>). *Journal of Biological Chemistry*. **279** (44): 45546–55. doi:10.1074/jbc.M402762200 (<https://doi.org/10.1074%2Fjbc.M402762200>). ISSN 0021-9258 (<https://www.worldcat.org/issn/0021-9258>). PMID 15326179 (<https://pubmed.ncbi.nlm.nih.gov/15326179>).
31. Brüssow H, Canchaya C, Hardt WD (September 2004). "Phages and the evolution of bacterial pathogens: from genomic rearrangements to lysogenic conversion" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC515249>). *Microbiology and Molecular Biology Reviews*. **68** (3): 560–602, table of contents. doi:10.1128/MMBR.68.3.560-602.2004 (<https://doi.org/10.1128%2FMMBR.68.3.560-602.2004>). PMC 515249 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC515249>). PMID 15353570 (<https://pubmed.ncbi.nlm.nih.gov/15353570>).
32. Krieg NR, Holt JG, eds. (1984). *Bergey's Manual of Systematic Bacteriology*. **1** (First ed.). Baltimore: The Williams & Wilkins Co. pp. 408–20. ISBN 978-0-683-04108-8.
33. Lukjancenko O, Wassenaar TM, Ussery DW (November 2010). "Comparison of 61 sequenced *Escherichia coli* genomes" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2974192>). *Microbial Ecology*. **60** (4): 708–20. doi:10.1007/s00248-010-9717-3 (<https://doi.org/10.1007%2Fs00248-010-9717-3>). PMC 2974192 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2974192>). PMID 20623278 (<https://pubmed.ncbi.nlm.nih.gov/20623278>).
34. Lan R, Reeves PR (September 2002). "Escherichia coli in disguise: molecular origins of Shigella". *Microbes and Infection*. **4** (11): 1125–32. doi:10.1016/S1286-4579(02)01637-4 (<https://doi.org/10.1016%2FS1286-4579%2802%2901637-4>). PMID 12361912 (<https://pubmed.ncbi.nlm.nih.gov/12361912>).
35. Orskov I, Orskov F, Jann B, Jann K (September 1977). "Serology, chemistry, and genetics of O and K antigens of *Escherichia coli*" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC414020>). *Bacteriological Reviews*. **41** (3): 667–710. doi:10.1128/MMBR.41.3.667-710.1977 (<https://doi.org/10.1128%2FMMBR.41.3.667-710.1977>). PMC 414020 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC414020>). PMID 334154 (<https://pubmed.ncbi.nlm.nih.gov/334154>).
36. Stenutz R, Weintraub A, Widmalm G (May 2006). "The structures of *Escherichia coli* O-polysaccharide antigens" (<https://doi.org/10.1111/j.1574-6976.2006.00016.x>). *FEMS Microbiology Reviews*. **30** (3): 382–403. doi:10.1111/j.1574-6976.2006.00016.x (<https://doi.org/10.1111%2Fj.1574-6976.2006.00016.x>). PMID 16594963 (<https://pubmed.ncbi.nlm.nih.gov/16594963>).
37. Lawrence JG, Ochman H (August 1998). "Molecular archaeology of the *Escherichia coli* genome" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC21352>). *Proceedings of the National Academy of Sciences of the United States of America*. **95** (16): 9413–17. Bibcode:1998PNAS...95.9413L (<https://ui.adsabs.harvard.edu/abs/1998PNAS...95.9413L>). doi:10.1073/pnas.95.16.9413 (<https://doi.org/10.1073%2Fpnas.95.16.9413>). PMC 21352 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC21352>). PMID 9689094 (<https://pubmed.ncbi.nlm.nih.gov/9689094>).
38. Nataro JP, Kaper JB (January 1998). "Diarrheagenic *Escherichia coli*" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC121379>). *Clinical Microbiology Reviews*. **11** (1): 142–201. doi:10.1128/CMR.11.1.142 (<https://doi.org/10.1128%2FCMR.11.1.142>). PMC 121379 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC121379>). PMID 9457432 (<https://pubmed.ncbi.nlm.nih.gov/9457432>).

39. Viljanen MK, Peltola T, Junnila SY, Olkkonen L, Järvinen H, Kuistila M, Huovinen P (October 1990). "Outbreak of diarrhoea due to *Escherichia coli* O111:B4 in schoolchildren and adults: association of Vi antigen-like reactivity". *Lancet*. **336** (8719): 831–34. doi:10.1016/0140-6736(90)92337-H (<https://doi.org/10.1016%2F0140-6736%2890%2992337-H>). PMID 1976876 (<https://pubmed.ncbi.nlm.nih.gov/1976876>). S2CID 23087850 (<https://api.semanticscholar.org/CorpusID:23087850>).
40. Battistuzzi FU, Feijao A, Hedges SB (November 2004). "A genomic timescale of prokaryote evolution: insights into the origin of methanogenesis, phototrophy, and the colonization of land" (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC533871>). *BMC Evolutionary Biology*. **4**: 44. doi:10.1186/1471-2148-4-44 (<https://doi.org/10.1186%2F1471-2148-4-44>). PMC 533871 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC533871>). PMID 15535883 (<https://pubmed.ncbi.nlm.nih.gov/15535883>).
41. Lecointre G, Rachdi L, Darlu P, Denamur E (December 1998). "Escherichia coli molecular phylogeny using the incongruence length difference test" (<https://doi.org/10.1093/oxfordjournals.molbev.a025895>). *Molecular Biology and Evolution*. **15** (12): 1685–95. doi:10.1093/oxfordjournals.molbev.a025895 (<https://doi.org/10.1093%2F0xfordjournals.molbev.a025895>). PMID 9866203 (<https://pubmed.ncbi.nlm.nih.gov/9866203>).
42. Bacteria make major evolutionary shift in the lab (<https://www.newscientist.com/channel/life/dn14094-bacteria-make-major-evolutionary-shift-in-the-lab.html>) *New Scientist*
43. Nair, Ramith R.; Vasse, Marie; Wielgoss, Sébastien; Sun, Lei; Yu, Yuen-Tsu N.; Velicer, Gregory J. "Bacterial predator-prey coevolution accelerates genome evolution and selects on virulence-associated prey defences", *Nature Communications*, 2019, 10:4301.
44. Brenner DJ, Krieg NR, Staley JT (26 July 2005) [1984 (Williams & Wilkins)]. Garrity GM (ed.). *The Gammaproteobacteria* (<https://www.springer.com/life+sciences/book/978-0-387-24144-9>). Bergey's Manual of Systematic Bacteriology. **2B** (2nd ed.). New York: Springer. p. 1108. ISBN 978-0-387-24144-9. British Library no. GBA561951.
45. Discussion of nomenclature of Enterobacteriaceae entry (<http://www.bacterio.net/e/enterobacteriaceae>) in LPSN [Euzéby, J.P. (1997). "List of Bacterial Names with Standing in Nomenclature: a folder available on the Internet" (<https://doi.org/10.1099/00207713-47-2-590>). *International Journal of Systematic and Evolutionary Microbiology*. **47** (2): 590–2. doi:10.1099/00207713-47-2-590 (<https://doi.org/10.1099%2F00207713-47-2-590>). PMID 9103655 (<https://pubmed.ncbi.nlm.nih.gov/9103655>).]
46. International Bulletin of Bacteriological Nomenclature and Taxonomy 8:73–74 (1958)
47. Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, Fiebig A, et al. (2013). "Complete genome sequence of DSM 30083(T), the type strain (U5/41(T)) of *Escherichia coli*, and a proposal for delineating subspecies in microbial taxonomy" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4334874>). *Standards in Genomic Sciences*. **9**: 2. doi:10.1186/1944-3277-9-2 (<https://doi.org/10.1186%2F1944-3277-9-2>). PMC 4334874 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4334874>). PMID 25780495 (<https://pubmed.ncbi.nlm.nih.gov/25780495>).
48. "Details: DSM-30083" (<http://www.dsmz.de/catalogues/details/culture/DSM-30083.html>). *dsmz.de*. Retrieved 10 January 2017.
49. "Escherichia coli (Migula) Castellani and Chalmers ATCC ® 11775&tra" (<http://www.atcc.org/ATCCAdvancedCatalogSearch/ProductDetails/tabid/452/Default.aspx?ATCCNum=11775&Template=bacteria>). *atcc.org*. Retrieved 10 January 2017.
50. "Escherichia" (<https://lpsn.dsmz.de/genus/escherichia>). LPSN. Retrieved 6 February 2011.
51. "Escherichia coli (Migula 1895) Castellani and Chalmers 1919" ([http://www.jcm.riken.go.jp/cgi-bin/jcm/jcm\\_number?JCM=1649](http://www.jcm.riken.go.jp/cgi-bin/jcm/jcm_number?JCM=1649)). *JCM Catalogue*.

52. Sims GE, Kim SH (May 2011). "Whole-genome phylogeny of *Escherichia coli*/Shigella group by feature frequency profiles (FFPs)" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3100984>). *Proceedings of the National Academy of Sciences of the United States of America*. **108** (20): 8329–34. Bibcode:2011PNAS..108.8329S (<https://ui.adsabs.harvard.edu/abs/2011PNAS..108.8329S>). doi:10.1073/pnas.1105168108 (<https://doi.org/10.1073%2Fpnas.1105168108>). PMC 3100984 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3100984>). PMID 21536867 (<https://pubmed.ncbi.nlm.nih.gov/21536867>).
53. Brzuszkiewicz E, Thürmer A, Schuldes J, Leimbach A, Liesegang H, Meyer FD, et al. (December 2011). "Genome sequence analyses of two isolates from the recent *Escherichia coli* outbreak in Germany reveal the emergence of a new pathotype: Entero-Aggregative-Haemorrhagic *Escherichia coli* (EAHEC)" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3219860>). *Archives of Microbiology*. **193** (12): 883–91. doi:10.1007/s00203-011-0725-6 (<https://doi.org/10.1007%2Fs00203-011-0725-6>). PMC 3219860 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3219860>). PMID 21713444 (<https://pubmed.ncbi.nlm.nih.gov/21713444>).
54. Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, et al. (September 1997). "The complete genome sequence of *Escherichia coli* K-12" (<https://doi.org/10.1126/science.277.5331.1453>). *Science*. **277** (5331): 1453–62. doi:10.1126/science.277.5331.1453 (<https://doi.org/10.1126%2Fscience.277.5331.1453>). PMID 9278503 (<https://pubmed.ncbi.nlm.nih.gov/9278503>).
55. Zhaxybayeva O, Doolittle WF (April 2011). "Lateral gene transfer". *Current Biology*. **21** (7): R242–46. doi:10.1016/j.cub.2011.01.045 (<https://doi.org/10.1016%2Fj.cub.2011.01.045>). PMID 21481756 (<https://pubmed.ncbi.nlm.nih.gov/21481756>). S2CID 14499247 (<https://api.semanticscholar.org/CorpusID:14499247>).
56. Hayashi K, Morooka N, Yamamoto Y, Fujita K, Isono K, Choi S, et al. (2006). "Highly accurate genome sequences of *Escherichia coli* K-12 strains MG1655 and W3110" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1681481>). *Molecular Systems Biology*. **2**: 2006.0007. doi:10.1038/msb4100049 (<https://doi.org/10.1038%2Fmsb4100049>). PMC 1681481 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1681481>). PMID 16738553 (<https://pubmed.ncbi.nlm.nih.gov/16738553>).
57. Zhou J, Rudd KE (January 2013). "EcoGene 3.0" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3531124>). *Nucleic Acids Research*. **41** (Database issue): D613–24. doi:10.1093/nar/gks1235 (<https://doi.org/10.1093%2Fnar%2Fgks1235>). PMC 3531124 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3531124>). PMID 23197660 (<https://pubmed.ncbi.nlm.nih.gov/23197660>).
58. Han MJ, Lee SY (June 2006). "The *Escherichia coli* proteome: past, present, and future prospects" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1489533>). *Microbiology and Molecular Biology Reviews*. **70** (2): 362–439. doi:10.1128/MMBR.00036-05 (<https://doi.org/10.1128%2FMMBR.00036-05>). PMC 1489533 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1489533>). PMID 16760308 (<https://pubmed.ncbi.nlm.nih.gov/16760308>).
59. Arifuzzaman M, Maeda M, Itoh A, Nishikata K, Takita C, Saito R, et al. (May 2006). "Large-scale identification of protein-protein interaction of *Escherichia coli* K-12" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1457052>). *Genome Research*. **16** (5): 686–91. doi:10.1101/gr.4527806 (<https://doi.org/10.1101%2Fgr.4527806>). PMC 1457052 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1457052>). PMID 16606699 (<https://pubmed.ncbi.nlm.nih.gov/16606699>).
60. Hu P, Janga SC, Babu M, Díaz-Mejía JJ, Butland G, Yang W, et al. (April 2009). Levchenko A (ed.). "Global functional atlas of *Escherichia coli* encompassing previously uncharacterized proteins" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2672614>). *PLOS Biology*. **7** (4): e96. doi:10.1371/journal.pbio.1000096 (<https://doi.org/10.1371%2Fjournal.pbio.1000096>). PMC 2672614 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2672614>). PMID 19402753 (<https://pubmed.ncbi.nlm.nih.gov/19402753>).
61. Rajagopala SV, Sikorski P, Kumar A, Mosca R, Vlasblom J, Arnold R, et al. (March 2014). "The binary protein-protein interaction landscape of *Escherichia coli*" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4123855>). *Nature Biotechnology*. **32** (3): 285–90. doi:10.1038/nbt.2831 (<https://doi.org/10.1038%2Fnbt.2831>). PMC 4123855 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4123855>). PMID 24561554 (<https://pubmed.ncbi.nlm.nih.gov/24561554>).

62. Todar K. "Pathogenic *E. coli*" (<http://www.textbookofbacteriology.net/e.coli.html>). *Online Textbook of Bacteriology*. University of Wisconsin–Madison Department of Bacteriology. Retrieved 30 November 2007.
63. Evans Jr DJ, Evans DG. "Escherichia Coli" (<https://web.archive.org/web/20071102062813/http://www.gsbs.utmb.edu/microbook/ch025.htm>). *Medical Microbiology, 4th edition*. The University of Texas Medical Branch at Galveston. Archived from the original (<http://www.gsbs.utmb.edu/microbook/ch025.htm>) on 2 November 2007. Retrieved 2 December 2007.
64. Wassenaar TM (September 2016). "E. Coli" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5063008>). *European Journal of Microbiology & Immunology*. **6** (3): 147–61. doi:10.1556/1886.2016.00029 (<https://doi.org/10.1556%2F1886.2016.00029>). PMC 5063008 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5063008>). PMID 27766164 (<https://pubmed.ncbi.nlm.nih.gov/27766164>).
65. Lodinová-Zádníková R, Cukrowska B, Tlaskalova-Hogenova H (July 2003). "Oral administration of probiotic Escherichia coli after birth reduces frequency of allergies and repeated infections later in life (after 10 and 20 years)". *International Archives of Allergy and Immunology*. **131** (3): 209–11. doi:10.1159/000071488 (<https://doi.org/10.1159%2F000071488>). PMID 12876412 (<https://pubmed.ncbi.nlm.nih.gov/12876412>). S2CID 19686481 (<https://api.semanticscholar.org/CorpusID:19686481>).
66. Grozdanov L, Raasch C, Schulze J, Sonnenborn U, Gottschalk G, Hacker J, Dobrindt U (August 2004). "Analysis of the genome structure of the nonpathogenic probiotic Escherichia coli strain Nissle 1917" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC490877>). *Journal of Bacteriology*. **186** (16): 5432–41. doi:10.1128/JB.186.16.5432-5441.2004 (<https://doi.org/10.1128%2FJB.186.16.5432-5441.2004>). PMC 490877 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC490877>). PMID 15292145 (<https://pubmed.ncbi.nlm.nih.gov/15292145>).
67. Kamada N, Inoue N, Hisamatsu T, Okamoto S, Matsuoka K, Sato T, et al. (May 2005). "Nonpathogenic Escherichia coli strain Nissle1917 prevents murine acute and chronic colitis". *Inflammatory Bowel Diseases*. **11** (5): 455–63. doi:10.1097/01.MIB.0000158158.55955.de (<https://doi.org/10.1097%2F01.MIB.0000158158.55955.de>). PMID 15867585 (<https://pubmed.ncbi.nlm.nih.gov/15867585>). S2CID 23386584 (<https://api.semanticscholar.org/CorpusID:23386584>).
68. "E. coli" (<http://www.mayoclinic.org/diseases-conditions/e-coli/basics/definition/con-20032105>). *mayoclinic.org – Mayo Clinic*. Retrieved 10 January 2017.
69. Lim JY, Yoon J, Hovde CJ (January 2010). "A brief overview of Escherichia coli O157:H7 and its plasmid O157" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3645889>). *Journal of Microbiology and Biotechnology*. **20** (1): 5–14. doi:10.4014/jmb.0908.08007 (<https://doi.org/10.4014%2Fjmb.0908.08007>). PMC 3645889 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3645889>). PMID 20134227 (<https://pubmed.ncbi.nlm.nih.gov/20134227>).
70. "E. coli" (<http://www.who.int/mediacentre/factsheets/fs125/en/>).
71. "E. coli Infection" (<https://www.cdc.gov/features/ecoliinfection/>). 15 June 2018.
72. "E. coli Food Poisoning." About. N.p., n.d. Web. 13 December 2014. <<http://www.about-ecoli.com/>>.
73. "Lung Congestion." TheFreeDictionary.com. N.p., n.d. Web. 13 December 2014. <<http://medical-dictionary.thefreedictionary.com/Lung+Congestion>>.
74. "Pulmonary Edema: Get the Facts on Treatment and Symptoms." MedicineNet. N.p., n.d. Web. 13 December 2014. <[http://www.medicinenet.com/pulmonary\\_edema/article.htm](http://www.medicinenet.com/pulmonary_edema/article.htm)>.
75. Staff, Mayo Clinic. "Hemolytic Uremic Syndrome (HUS)." Mayo Clinic. Mayo Foundation for Medical Education and Research, 3 July 2013. Web. 13 December 2014. <<http://www.mayoclinic.com/health/hemolytic-uremic-syndrome/DS00876>>.
76. "Uropathogenic Escherichia coli: The Pre-Eminent Urinary Tract Infection Pathogen" ([https://web.archive.org/web/20131202232432/https://www.novapublishers.com/catalog/product\\_info.php?products\\_id=25500&osCsid=3712df5600f98259a8bdc1d9baf202e9](https://web.archive.org/web/20131202232432/https://www.novapublishers.com/catalog/product_info.php?products_id=25500&osCsid=3712df5600f98259a8bdc1d9baf202e9)). Nova publishers. Archived from the original ([https://www.novapublishers.com/catalog/product\\_info.php?products\\_id=25500&osCsid=3712df5600f98259a8bdc1d9baf202e9](https://www.novapublishers.com/catalog/product_info.php?products_id=25500&osCsid=3712df5600f98259a8bdc1d9baf202e9)) on 2 December 2013. Retrieved 27 November 2013.



77. Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB (October 2013). "Recent advances in understanding enteric pathogenic *Escherichia coli*" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3811233>). *Clinical Microbiology Reviews*. **26** (4): 822–80. doi:10.1128/CMR.00022-13 (<https://doi.org/10.1128/CMR.00022-13>). PMC 3811233 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3811233>). PMID 24092857 (<https://pubmed.ncbi.nlm.nih.gov/24092857>).
78. "Outbreaks of *E. coli* O104:H4 infection: update 29" (<https://web.archive.org/web/20110808155129/http://www.euro.who.int/en/what-we-do/health-topics/emergencies/international-health-regulations/news/news/2011/07/outbreaks-of-e.-coli-o104h4-infection-update-29>). WHO. 7 July 2011. Archived from the original (<http://www.euro.who.int/en/what-we-do/health-topics/emergencies/international-health-regulations/news/news/2011/07/outbreaks-of-e.-coli-o104h4-infection-update-29>) on 8 August 2011.
79. "Samen von Bockshornklee mit hoher Wahrscheinlichkeit für EHEC O104:H4 Ausbruch verantwortlich in English: *Fenugreek seeds with high probability for EHEC O104: H4 responsible outbreak*" ([http://www.bfr.bund.de/cm/343/samen\\_von\\_bockshornklee\\_mit\\_hoher\\_wahrscheinlichkeit\\_fuer\\_ehec\\_o104\\_h4\\_ausbruch\\_verantwortlich.pdf](http://www.bfr.bund.de/cm/343/samen_von_bockshornklee_mit_hoher_wahrscheinlichkeit_fuer_ehec_o104_h4_ausbruch_verantwortlich.pdf)) (PDF) (in German). Bundesinstitut für Risikobewertung (BfR) (Federal Institute for Risk Assessment). 30 June 2011. Retrieved 17 July 2011.
80. "General Information| *E.coli* | CDC" (<https://www.cdc.gov/ecoli/general/>). *www.cdc.gov*. Retrieved 19 April 2017.
81. US Centers for Disease Control and Prevention. "Enterotoxigenic *E. coli* (ETEC)" (<https://www.cdc.gov/ecoli/etec.html>). Retrieved 21 July 2016.
82. Al-Abri SS, Beeching NJ, Nye FJ (June 2005). "Traveller's diarrhoea". *The Lancet. Infectious Diseases*. **5** (6): 349–60. doi:10.1016/S1473-3099(05)70139-0 ([https://doi.org/10.1016/S1473-3099\(05\)70139-0](https://doi.org/10.1016/S1473-3099(05)70139-0)). PMID 15919621 (<https://pubmed.ncbi.nlm.nih.gov/15919621>).
83. Bourgeois AL, Wierzba TF, Walker RI (June 2016). "Status of vaccine research and development for enterotoxigenic *Escherichia coli*" (<https://doi.org/10.1016/j.vaccine.2016.02.076>). *Vaccine*. **34** (26): 2880–86. doi:10.1016/j.vaccine.2016.02.076 (<https://doi.org/10.1016/j.vaccine.2016.02.076>). PMID 26988259 (<https://pubmed.ncbi.nlm.nih.gov/26988259>).
84. Svennerholm AM (February 2011). "From cholera to enterotoxigenic *Escherichia coli* (ETEC) vaccine development" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3089050>). *The Indian Journal of Medical Research*. **133**: 188–96. PMC 3089050 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3089050>). PMID 21415493 (<https://pubmed.ncbi.nlm.nih.gov/21415493>).
85. Farrar J, Hotez P, Junghanss T, Kang G, Lalloo D, White NJ, eds. (2013). *Manson's Tropical Diseases* (23rd ed.). Oxford: Elsevier/Saunders. ISBN 978-0702053061.
86. "General Information- *E.coli*" (<https://www.cdc.gov/ecoli/general/index.html>). Centers for Disease Control and Prevention. Retrieved 25 May 2017.
87. Lee SY (March 1996). "High cell-density culture of *Escherichia coli*". *Trends in Biotechnology*. **14** (3): 98–105. doi:10.1016/0167-7799(96)80930-9 ([https://doi.org/10.1016/0167-7799\(96\)80930-9](https://doi.org/10.1016/0167-7799(96)80930-9)). PMID 8867291 (<https://pubmed.ncbi.nlm.nih.gov/8867291>).
88. Russo E (January 2003). "The birth of biotechnology". *Nature*. **421** (6921): 456–57. Bibcode:2003Natur.421..456R (<https://ui.adsabs.harvard.edu/abs/2003Natur.421..456R>). doi:10.1038/nj6921-456a (<https://doi.org/10.1038/nj6921-456a>). PMID 12540923 (<https://pubmed.ncbi.nlm.nih.gov/12540923>). S2CID 4357773 (<https://api.semanticscholar.org/CorpusID:4357773>).
89. Cornelis P (October 2000). "Expressing genes in different *Escherichia coli* compartments". *Current Opinion in Biotechnology*. **11** (5): 450–54. doi:10.1016/S0958-1669(00)00131-2 ([https://doi.org/10.1016/S0958-1669\(00\)00131-2](https://doi.org/10.1016/S0958-1669(00)00131-2)). PMID 11024362 (<https://pubmed.ncbi.nlm.nih.gov/11024362>).
90. Tof I (1994). "Recombinant DNA Technology in the Synthesis of Human Insulin" (<http://www.littletree.com.au/dna.htm>). Little Tree Pty. Ltd. Retrieved 30 November 2007.

91. Bessette PH, Aslund F, Beckwith J, Georgiou G (November 1999). "Efficient folding of proteins with multiple disulfide bonds in the Escherichia coli cytoplasm" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC24128>). *Proceedings of the National Academy of Sciences of the United States of America*. **96** (24): 13703–08. Bibcode:1999PNAS...9613703B (<https://ui.adsabs.harvard.edu/abs/1999PNAS...9613703B>). doi:10.1073/pnas.96.24.13703 (<https://doi.org/10.1073/pnas.96.24.13703>). PMC 24128 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC24128>). PMID 10570136 (<https://pubmed.ncbi.nlm.nih.gov/10570136>).
92. Ihssen J, Kowarik M, Dilettoso S, Tanner C, Wacker M, Thöny-Meyer L (August 2010). "Production of glycoprotein vaccines in Escherichia coli" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2927510>). *Microbial Cell Factories*. **9** (61): 61. doi:10.1186/1475-2859-9-61 (<https://doi.org/10.1186/1475-2859-9-61>). PMC 2927510 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2927510>). PMID 20701771 (<https://pubmed.ncbi.nlm.nih.gov/20701771>).
93. Wacker M, Linton D, Hitchen PG, Nita-Lazar M, Haslam SM, North SJ, et al. (November 2002). "N-linked glycosylation in Campylobacter jejuni and its functional transfer into E. coli". *Science*. **298** (5599): 1790–93. Bibcode:2002Sci...298.1790W (<https://ui.adsabs.harvard.edu/abs/2002Sci...298.1790W>). doi:10.1126/science.298.5599.1790 (<https://doi.org/10.1126/science.298.5599.1790>). PMID 12459590 (<https://pubmed.ncbi.nlm.nih.gov/12459590>).
94. Huang CJ, Lin H, Yang X (March 2012). "Industrial production of recombinant therapeutics in Escherichia coli and its recent advancements". *Journal of Industrial Microbiology & Biotechnology*. **39** (3): 383–99. doi:10.1007/s10295-011-1082-9 (<https://doi.org/10.1007/s10295-011-1082-9>). PMID 22252444 (<https://pubmed.ncbi.nlm.nih.gov/22252444>). S2CID 15584320 (<https://api.semanticscholar.org/CorpusID:15584320>).
95. Summers, Rebecca (24 April 2013) Bacteria churn out first ever petrol-like biofuel (<https://www.newscientist.com/article/dn23431-bacteria-churn-out-first-ever-petrol-like-biofuel.html>) New Scientist, Retrieved 27 April 2013
96. Nic Halverson (15 August 2013). "Bacteria-Powered Light Bulb Is Electricity-Free" (<http://news.discovery.com/tech/alternative-power-sources/bacteria-powered-light-bulb-is-electricity-free-130815.htm>).
97. Ninfa AJ, Ballou DP (2009). *Fundamental Laboratory Approaches for Biochemistry and Biotechnology*. Wiley. p. 230. ISBN 978-0470087664.
98. "Cloning, sequence analysis, and expression of cDNA coding for the major house dust mite allergen, Der f 1, in Escherichia coli" *Brazilian Journal of Medical and Biological Research*
99. Fux CA, Shirtliff M, Stoodley P, Costerton JW (February 2005). "Can laboratory reference strains mirror "real-world" pathogenesis?". *Trends in Microbiology*. **13** (2): 58–63. doi:10.1016/j.tim.2004.11.001 (<https://doi.org/10.1016/j.tim.2004.11.001>). PMID 15680764 (<https://pubmed.ncbi.nlm.nih.gov/15680764>).
00. Vidal O, Longin R, Prigent-Combaret C, Dorel C, Hooreman M, Lejeune P (May 1998). "Isolation of an Escherichia coli K-12 mutant strain able to form biofilms on inert surfaces: involvement of a new ompR allele that increases curli expression" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC107187>). *Journal of Bacteriology*. **180** (9): 2442–49. doi:10.1128/JB.180.9.2442-2449.1998 (<https://doi.org/10.1128/JB.180.9.2442-2449.1998>). PMC 107187 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC107187>). PMID 9573197 (<https://pubmed.ncbi.nlm.nih.gov/9573197>).
01. TiO<sub>2</sub> coatings as bactericidal photocatalysts in water treatment ([https://www.researchgate.net/publication/235903481\\_The\\_effects\\_of\\_firing\\_conditions\\_on\\_the\\_properties\\_of\\_electrophoretically\\_deposited\\_titanium\\_dioxide\\_films\\_on\\_graphite\\_substrates](https://www.researchgate.net/publication/235903481_The_effects_of_firing_conditions_on_the_properties_of_electrophoretically_deposited_titanium_dioxide_films_on_graphite_substrates)).
02. Lederberg J, Tatum EL (October 1946). "Gene recombination in Escherichia coli" ([http://profiles.nlm.nih.gov/BB/G/A/S/Z/\\_/bbgasz.pdf](http://profiles.nlm.nih.gov/BB/G/A/S/Z/_/bbgasz.pdf)) (PDF). *Nature*. **158** (4016): 558. Bibcode:1946Natur.158..558L (<https://ui.adsabs.harvard.edu/abs/1946Natur.158..558L>). doi:10.1038/158558a0 (<https://doi.org/10.1038/158558a0>). PMID 21001945 (<https://pubmed.ncbi.nlm.nih.gov/21001945>). S2CID 1826960 (<https://api.semanticscholar.org/CorpusID:1826960>). Source: National Library of Medicine – The Joshua Lederberg Papers (<http://profiles.nlm.nih.gov/BB/G/A/S/Z/>)
03. *Biological Activity of Crystal*. p. 169.

04. Susman M (March 1995). "The Cold Spring Harbor Phage Course (1945–1970): a 50th anniversary remembrance" (<https://web.archive.org/web/20060916155323/https://www.cshl.edu/History/phagecourse.html>). *Genetics*. **139** (3): 1101–06. PMC [1206443](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1206443) (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1206443>). PMID [7768426](https://pubmed.ncbi.nlm.nih.gov/7768426) (<https://pubmed.ncbi.nlm.nih.gov/7768426>). Archived from the original (<https://www.cshl.edu/History/phagecourse.html>) on 16 September 2006.
05. Benzer S (March 1961). "On the Topography of the Genetic Fine Structure" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC221592>). *Proceedings of the National Academy of Sciences of the United States of America*. **47** (3): 403–15. Bibcode:1961PNAS...47..403B (<https://ui.adsabs.harvard.edu/abs/1961PNAS...47..403B>). doi:10.1073/pnas.47.3.403 (<https://doi.org/10.1073/pnas.47.3.403>). PMC [221592](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC221592) (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC221592>). PMID [16590840](https://pubmed.ncbi.nlm.nih.gov/16590840) (<https://pubmed.ncbi.nlm.nih.gov/16590840>).
06. "Facts about *E. Coli*" (<http://eol.org/pages/972688/details>). Encyclopedia of Life. Retrieved 27 November 2013.
07. "Scarab Genomics LLC. Company web site" (<http://www.scarabgenomics.com/>).
08. Umenhoffer K, Fehér T, Balikó G, Ayaydin F, Pósfai J, Blattner FR, Pósfai G (May 2010). "Reduced evolvability of *Escherichia coli* MDS42, an IS-less cellular chassis for molecular and synthetic biology applications" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2891674>). *Microbial Cell Factories*. **9**: 38. doi:10.1186/1475-2859-9-38 (<https://doi.org/10.1186/1475-2859-9-38>). PMC [2891674](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2891674) (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2891674>). PMID [20492662](https://pubmed.ncbi.nlm.nih.gov/20492662) (<https://pubmed.ncbi.nlm.nih.gov/20492662>).
09. Pósfai G, Plunkett G, Fehér T, Frisch D, Keil GM, Umenhoffer K, et al. (May 2006). "Emergent properties of reduced-genome *Escherichia coli*". *Science*. **312** (5776): 1044–46. Bibcode:2006Sci...312.1044P (<https://ui.adsabs.harvard.edu/abs/2006Sci...312.1044P>). doi:10.1126/science.1126439 (<https://doi.org/10.1126/science.1126439>). PMID [16645050](https://pubmed.ncbi.nlm.nih.gov/16645050) (<https://pubmed.ncbi.nlm.nih.gov/16645050>). S2CID [43287314](https://api.semanticscholar.org/CorpusID:43287314) (<https://api.semanticscholar.org/CorpusID:43287314>).
10. Kolisnychenko V, Plunkett G, Herring CD, Fehér T, Pósfai J, Blattner FR, Pósfai G (April 2002). "Engineering a reduced *Escherichia coli* genome" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC187512>). *Genome Research*. **12** (4): 640–47. doi:10.1101/gr.217202 (<https://doi.org/10.1101/gr.217202>). PMC [187512](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC187512) (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC187512>). PMID [11932248](https://pubmed.ncbi.nlm.nih.gov/11932248) (<https://pubmed.ncbi.nlm.nih.gov/11932248>).
11. Keymer JE, Galajda P, Muldoon C, Park S, Austin RH (November 2006). "Bacterial metapopulations in nanofabricated landscapes" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1635019>). *Proceedings of the National Academy of Sciences of the United States of America*. **103** (46): 17290–95. Bibcode:2006PNAS..10317290K (<https://ui.adsabs.harvard.edu/abs/2006PNAS..10317290K>). doi:10.1073/pnas.0607971103 (<https://doi.org/10.1073/pnas.0607971103>). PMC [1635019](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1635019) (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1635019>). PMID [17090676](https://pubmed.ncbi.nlm.nih.gov/17090676) (<https://pubmed.ncbi.nlm.nih.gov/17090676>).
12. Baumgardner J, Acker K, Adefuye O, Crowley ST, Deloache W, Dickson JO, et al. (July 2009). "Solving a Hamiltonian Path Problem with a bacterial computer" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2723075>). *Journal of Biological Engineering*. **3**: 11. doi:10.1186/1754-1611-3-11 (<https://doi.org/10.1186/1754-1611-3-11>). PMC [2723075](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2723075) (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2723075>). PMID [19630940](https://pubmed.ncbi.nlm.nih.gov/19630940) (<https://pubmed.ncbi.nlm.nih.gov/19630940>).
13. Tirumalai MR, Karouia F, Tran Q, Stepanov VG, Bruce RJ, Ott M, Pierson DL, Fox GE (May 2017). "The adaptation of *Escherichia coli* cells grown in simulated microgravity for an extended period is both phenotypic and genomic" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5460176>). *npj Microgravity*. **3** (15). doi:10.1038/s41526-017-0020-1 (<https://doi.org/10.1038/s41526-017-0020-1>). PMC [5460176](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5460176) (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5460176>). PMID [28649637](https://pubmed.ncbi.nlm.nih.gov/28649637) (<https://pubmed.ncbi.nlm.nih.gov/28649637>).

14. Tirumalai MR, Karouia F, Tran Q, Stepanov VG, Bruce RJ, Ott M, Pierson DL, Fox GE (January 2019). "Evaluation of acquired antibiotic resistance in *Escherichia coli* exposed to long-term low-shear modeled microgravity and background antibiotic exposure" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6336426>). *mBio*. **10** (e02637-18). doi:10.1128/mBio.02637-18 (<https://doi.org/10.1128%2FmBio.02637-18>). PMC 6336426 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6336426>). PMID 30647159 (<https://pubmed.ncbi.nlm.nih.gov/30647159>).
15. Haeckel, Ernst (1867). *Generelle Morphologie der Organismen*. Reimer, Berlin. ISBN 978-1-144-00186-3.
16. Escherich T (1885). "Die Darmbakterien des Neugeborenen und Säuglinge" (<https://books.google.com/books?id=o1MXAAAAYAAJ&lpg=PA135&dq=%22Die%20darmbakterien%20des%20neugeborene%20und%20säuglings%22&pg=PA135#v=onepage>). *Fortschr. Med.* **3**: 515–22.
17. Breed RS, Conn HJ (May 1936). "The Status of the Generic Term Bacterium Ehrenberg 1828" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC543738>). *Journal of Bacteriology*. **31** (5): 517–18. doi:10.1128/JB.31.5.517-518.1936 (<https://doi.org/10.1128%2FJB.31.5.517-518.1936>). PMC 543738 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC543738>). PMID 16559906 (<https://pubmed.ncbi.nlm.nih.gov/16559906>).
18. Migula W (1895). "Bacteriaceae (Stäbchenbakterien)". In Engerl A, Prantl K (eds.). *Die Natürlichen Pflanzenfamilien, W. Engelmann, Leipzig, Teil I, Abteilung Ia*. pp. 20–30.
19. Castellani A, Chalmers AJ (1919). *Manual of Tropical Medicine* (<https://archive.org/details/manualoftrropical00cast>) (3rd ed.). New York: Williams Wood and Co.
20. "BBC News | Health | Sheriff criticises E. Coli butcher" (<http://news.bbc.co.uk/1/hi/health/154107.stm>).

## Databases and external links

---

- **EcoCyc** – literature-based curation of the entire genome, and of transcriptional regulation, transporters, and metabolic pathways
- **Membranome database** provides information about single-pass transmembrane proteins from *E. coli* and several other organisms
- *E. coli* statistics ([http://redpoll.pharmacy.ualberta.ca/CCDB/cgi-bin/STAT\\_NEW.cgi](http://redpoll.pharmacy.ualberta.ca/CCDB/cgi-bin/STAT_NEW.cgi))
- *E. coli* Infection | Causes & Risk Factors (<http://familydoctor.org/familydoctor/en/diseases-conditions/ecoli-infection/causes-risk-factors.html>)
- **Bacteriome** (<http://www.compsysbio.org/bacteriome/>) *E. coli* interaction database
- **EcoGene** (<https://web.archive.org/web/20180523221348/http://www.ecogene.org/>) (genome database and website dedicated to *Escherichia coli* K-12 substrain MG1655)
- **EcoSal** (<https://web.archive.org/web/20131011071918/http://ecosal.org/>) Continually updated Web resource based on the classic ASM Press publication *Escherichia coli and Salmonella: Cellular and Molecular Biology*
- **ECODAB** (<https://web.archive.org/web/20140609195946/http://www.casper.organ.su.se/ECODAB/>) The structure of the O-antigens that form the basis of the serological classification of *E. coli*
- **Coli Genetic Stock Center** (<http://cgsc.biology.yale.edu/index.php>) Strains and genetic information on *E. coli* K-12
- **PortEco** (formerly EcoliHub) (<http://porteco.org>) – NIH-funded comprehensive data resource for *E. coli* K-12 and its phage, plasmids, and mobile genetic elements
- **EcoliWiki** (<http://ecoliwiki.net>) is the community annotation component of PortEco (<http://porteco.org>)
- **RegulonDB** (<http://regulondb.ccg.unam.mx/index.jsp>) RegulonDB is a model of the complex regulation of transcription initiation or regulatory network of the cell *E. coli* K-12.
- **Uropathogenic Escherichia coli (UPEC)** (<http://www.genome.wisc.edu/sequencing/upec.htm>)

- [AlignACE \(http://arep.med.harvard.edu/ecoli\\_matrices/\)](http://arep.med.harvard.edu/ecoli_matrices/) Matrices that search for additional binding sites in the *E. coli* genomic sequence
- 

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Escherichia\\_coli&oldid=978717330](https://en.wikipedia.org/w/index.php?title=Escherichia_coli&oldid=978717330)"

---

**This page was last edited on 16 September 2020, at 15:09 (UTC).**

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.